

Hibernal Diapause of North American Papilionoidea and Hesperioidea

James A. Scott

60 Estes Street, Lakewood, Colorado 80226

Abstract. Information from 695 sources (mainly scientific journals) is presented on the life history stage in which 348 species of butterflies and skippers spend the winter in North America. *Parnassius*, *Neophasia*, many Theclini (especially *Satyrium*), most *Lycaena*, and a few others diapause as eggs. Coliadinae, Satyrinae, most Nymphalidae, Riodininae, two *Lycaena* of Palearctic affinity, many Plebejini, and most Hesperioidea diapause as larvae. Most Papilionoidea, Pierinae, many Theclini (especially *Callophrys*), many Polyommata, and a few others diapause as pupae. *Danaus plexippus*, *Anaea andria*, *Nymphalis*, *Polygonia*, and some *Vanessa* diapause as adults. Records from Japan and Britain are very similar. The boreal or arctic-alpine species which have a two-year life cycle generally pass the first winter as a young larva, the second winter as a mature larva. The life history stage in which winter is passed significantly influences the time of flight of adult broods during the year.

Introduction

Insects in temperate climates must survive the cold temperatures of winter. To do this, insects have a phenomenon called hibernal diapause. Diapause is an arrested state of development. Growth and feeding of larvae stop, embryonic and pupal development cease, and mating and reproduction (egg development) of adults do not occur. Diapause is usually noticed when rearing is attempted; when the immatures reach a certain stage, they do not feed or develop further, and unless special precautions are taken, they will desiccate and die. The basic function of diapause is to allow insects to endure an unfavorable time such as a freezing winter. The tropical species which stray into temperate climates usually do not have diapause stages, and they soon die in temperate zone winters. The native temperate zone species, however, spend the winter in a diapause stage. Insects in general can spend the winter in any stage, including egg, larva, pupa, or adult, but a given species is usually adapted to diapause in only one stage. A few species may be able to winter in several stages, however. Some alpine and arctic species which have a two-year life cycle may winter in different stages during the two winters, as noted below. Good general discussions of diapause are by Chapman (2), Danilevskii (4), Beck (1), and Lees (14).

Physiological and sometimes morphological changes occur during diapause. Growth stops during diapause. During winter diapause insects increase the concentration of glycerol, sorbitol, or alcohol in their bodies, which acts as antifreeze to lower the temperature at which they freeze. Free water is converted to a colloid-bound state, preventing damage from freezing (4). Insects in diapause respire at a rate much less than normal (454), and they are quiescent and do not feed or grow. Diapausing larvae usually have one extra instar, which is usually specialized for winter survival (132, 459). The hibernating generation of *Asterocampa* has six instars, versus 5 for the summer generation, for example. *Chlosyne gorgone* and *nycteis* have a special diapause instar with russet colored larvae (529, 97). *Plebejus icarioides* larvae turn brown in the second instar when they diapause, and are green before and after that (535, 536). *Euphydryas editha* has a special diapause instar with a thicker integument (548).

Diapause of adult butterflies such as *Nymphalis* takes the form of an arrested state of development of their reproductive system; the abdominal fat in females is not converted to eggs until diapause is broken.

Two environmental factors, sunlight and temperature, regulate the appearance of diapause in most insects (1, 2, 4, 14). The number of hours of sunlight during the day (the photoperiod) has been found to be the critical factor in starting diapause in most insects. Diapause occurs only when photoperiod is below a critical amount. One stage in the life cycle is sensitive to photoperiod, which causes that stage or (usually) a later one to undergo diapause. Egg diapause is usually initiated by short photoperiod experienced by females laying the eggs, larval diapause by short photoperiod experienced by an earlier larval stage, pupal diapause by short photoperiod experienced by the larva, and adult diapause by short photoperiod experienced in larvae or (in ladybird beetles) the adult. For example, *Limenitis archippus* diapauses as larvae (306), *Papilio polyxenes* as pupae (305), and *Pieris napi* as pupae (377). In all three species, the long days of spring and early summer prevent diapause, but the short days of late summer and fall act on the larva to trigger diapause. In some insects high temperatures prevent the onset of diapause and low temperatures may bring on diapause. High temperatures prevent diapause in *Lycaena phlaeas* (334). In *Pieris rapae*, diapause never occurs at 24 degrees C, but at 20 degrees C, the photoperiod experienced by the fourth-instar larva influences the pupal diapause; with 10-hour days all the pupae diapause, but with 13-hour days none diapause (289).

In many species including *Limenitis archippus*, *Pieris rapae* and *P. napi* the photoperiod mechanism differs between geographic locations (306c, 1, 14); the critical photoperiod allowing continued development increases with latitude, because the diapause mechanism has adapted to the local sunlight conditions at each latitude.

Many species with only one brood per year have obligatory diapause; when a particular stage is reached, diapause occurs regardless of normal environmental conditions. Examples are *Papilio indra fordi*, *Anthocharis cethura* (535), *Euphydryas phaeton* (393a), and many *Satyrium*. Physiologically, this usually occurs because the critical photoperiod allowing development is greater than that found at the latitude of the habitat. Extreme laboratory conditions (constant light and warmth) sometimes can prevent diapause in these species (4). Ecologically, obligatory diapause is generally due to the absence of food plants or other necessities later in the season.

Generally, prolonged exposure to cold in the winter is necessary to "break" diapause and allow the insects to resume growing when the weather warms. Long days also break diapause in *P. polyxenes* (305). Diapause may terminate spontaneously merely due to the passage of time (40a).

The net effect of these mechanisms is that development usually starts when the weather warms in spring, and ceases when photoperiod triggers diapause (most multivoltine species), or when the diapause stage is reached (in the obligatory-diapause univoltine species).

This paper discusses only winter (hibernal) diapause, but summer (aestival) diapause sometimes occurs as well (328). The same cessation of development occurs in aestival diapause as in hibernal diapause, but it usually occurs during a time of year when droughts are frequent. Adult diapause is difficult to recognize; it is suspected when lifespan is unusually long and mating and reproduction do not occur, and dissections of females reveal fat body but no mature eggs. *Coenonympha tullia* adults (40) and *Euphydryas* larvae (509) may diapause in the summer in California which has a mediterranean climate. *Ochlodes sylvanoides* has a short mature larval diapause in early summer in California (535). *Agathymus* larvae may aestivate (12). *Speyeria* species have only one generation per year, but the flight period is very long (usually mid June to mid September), and diapause seems to occur during it. Edwards (5) and Scudder (21) found that *S. cybele*, *aphrodite*, and *idalia* females do not lay eggs until late August or September. Scudder found that *S. cybele* eggs do not become full size inside the female until mid August, and in Virginia adults are rarely seen in July (they are apparently diapausing). *Speyeria coronis* and *zereke* also have an adult diapause in summer (296a). The physiological factors producing aestival diapause are not as well known as those for hibernal diapause. In general long days and high temperatures cause aestival diapause. A geometrid moth *Abraxes miranda* has a short aestival diapause caused by long days and moderate temperatures (1). Low temperatures cannot play a large role in breaking aestival diapause, because summer is warm.

Results

This section presents the available data on hibernal diapause stage. The data are arranged in several tables. Table 1 contains the mostly temperate zone species which almost certainly have evolved a diapause stage. Table 2 contains the species which probably do not have a true diapause. These are semitropical species or species which may migrate into temperate regions each year. These species are often reported to spend the winter in several different stages; species without diapause continue to develop so that all stages can be present at a given time whenever the weather is mild enough not to kill them.

These data are based on 623 literature sources, on personal communications from 27 scientists, and on 46 of my own records. I searched all of the major entomology journals which have many papers on butterflies, and nearly all of the major works and papers on butterflies. Some references have undoubtedly been missed, but it is significant that the last several hundred references examined produced very few species-stage records not already found. Because slightly more than half of the approximately 700 species found in North America lack data on diapause stage, future contributions to the subject will come from rearing this unknown fraction and the species inadequately studied before now.

The data are cited by number (see Literature Cited). Authors of scientific names are not listed; they can be found in the references cited. The taxonomy of (427, 13, 12) is used except for recent changes that will appear in my forthcoming "Field Guide to North American Butterflies."

Discussion and Conclusions

Table 3 summarizes winter diapause information for nearctic species with adequate data, and compares it to Japanese (348) and British (11) information. Results are very similar among all three faunas. *Parnassius* usually diapause as eggs, but the remaining Papilionidae diapause as pupae. In the Pieridae, the Coliadinae usually diapause as larvae whereas the Pierinae usually diapause as pupae. Among the Nymphalidae, the Satyrinae and most other Nymphalidae almost always diapause as larvae, although *Anaea andria*, and the genera *Polygonia*, *Nymphalis*, *Aglais*, and some *Vanessa* (these four are closely related) diapause as adults. Among the Lycaenidae, the Theclini diapause as eggs usually, except for a large section (mainly *Callophrys*) that diapauses as pupae. Most American *Lycaena* diapause as eggs, but two American species (*phlaeas* and *cupreus*) belong to a Palearctic group which diapauses as larvae. The Polyommatini usually diapause as larvae, occasionally as eggs or pupae. The Riodininae studied diapause as larvae. Hesperidae usually diapause as larvae.

The larval diapause species are usually specialized to diapause as young, part-grown, or mature larvae. Most diapause as part-grown larvae, but a

few (*Cercyonis*, *Speyeria*, and others) diapause as first instar larvae, and *Everes*, most *Pyrginae*, and several others diapause as mature larvae.

Several-year diapause of annual species. Occasionally species with life cycles of one year or less have longer life cycles because diapausing individuals do not emerge in spring but emerge only after another year or more in diapause. *Euphydryas chalcedona* larvae sometimes diapause for up to five years (54), and *E. editha* sometimes diapause for 2 years as larvae (138). Many pupal diapause species are known to do this (5, 385), including *Papilio aristodemus* (13, 377b), *P. cresphontes* (21, 95), *P. bairdii* (170), *P. zelicaon* (538, 315a, 335a), *P. polyxenes* (234, 101), *P. p. rudkini* (503, 477a, 22a, 427--six years), *P. indra* (412), *P. multicaudata* (167), *P. glaucus* (234), *P. g. rutulus* (167), *Eurytides marcellus* (234), *Anthocharis sara* (471, 167, 343a, 387a, 477a), *A. cethura* (471, 477a, 167, 535--five years), *A. lanceolata* (254, 477a, 421), *A. midea* (167, 234), *Euchloe ausonides* (167, 477a), *E. hyantis* (477a), *Pieris rapae* (482), and *Pieris sisymbri* (477a, 535--five years). Eggs might also diapause for several years. Several-year diapause would seem to be a useful adaptation to avoiding unfavorable years (290). Emergence of *Papilio aristodemus* (377b), *Anthocharis cethura* and *Euchloe hyantis* (535), seems to be stimulated by moisture in addition to cold. *Papilio polyxenes rudkini* and *Chlosyne neumoegeni* have summer and fall broods that seem to emerge from diapause as a response to rain; the more the rain, the larger the emergence (535).

Diapause stage of biennial species. Some boreal, arctic, and alpine species always have a two-year life cycle. They fly as adults only on alternate years, or at a given locality there are even-year and odd-year cohorts which rarely or never interbreed. Table 4 lists the species which are biennial, and summarizes the data on diapause stages from Table 1 for these species. For *Parnassius evermanni* diapause occurs as an egg the first winter, then pupa the second winter, so that larval feeding occurs only on alternate years. For most other species, however, which include two *Colias*, *Chlosyne damoetas* (probably biennial), *Boloria alberta* ?, and seven *Oeneis*, the young larva spends the first winter, and the older or mature larva spends the next winter. This interpretation nicely explains the data on winter stages of *Oeneis*, which otherwise seem contradictory or confusing. Many other species of *Erebia*, *Boloria*, etc., are suspected of being biennial.

One very interesting aspect of biennialism is that when several biennial species occur together, they often fly on different years. For instance, *Oeneis chryxus* and *Neominois ridingsii* fly on Sonora Pass, Mono Co., California, at the same sites, *chryxus* in odd years, *ridingsii* in even years. *Oeneis jutta* and *O. chryxus* alternate years in Michigan, *O. jutta* and *O. (nevadensis?) macounii* alternate years in Minnesota. Adult interference during mate-locating behavior or larval competition may cause these species to fly in alternate years. Larval feeding in biennial species may be

concentrated or restricted to alternate years. In eastern Alaska around Fairbanks, however, *O. jutta*, *O. polixenes*, *Erebia disa*, *Boloria astarte distincta*, *B. polaris*, *B. (titania?) chariclea*, and *Hesperia comma* all fly in odd-numbered years. In such cases perhaps the weather was so unfavorable in the past in an even-numbered year that all the adult cohort failed to mate and reproduce, and only the cohort occurring on the odd-years survived. Collectors returning from the far north often report such weather. Eventually, when more of the cohorts for biennial species have been discovered and mapped, a comparison with Pleistocene climatic events may help elucidate these patterns.

Non-diapause species. Contradictory information sometimes occurs for subtropical species which have no definite diapause stage (Table 2). Since these species develop during the winter, they may be found then in any stage. For instance, *Ascia monuste* develops year-around in Florida, and freezing winters kill the entire population (47). Some species such as *Euptoieta claudia* and *Precis coenia*, which have been reported to spend the winter in several stages, are warm-weather species which may actually have no diapause stage. Several species (*Colias eurytheme*, *Vanessa cardui*) may or may not have a true diapause. Hovanitz (525) and Shapiro (318) think that *eurytheme* may not have a true diapause, but the best data for this species indicate that it passes the cold winters in the larval stage as do its congeners with a true diapause. It has expanded its range northward within the last 100 years and may not have adapted yet to northern latitude photoperiods. Its close relative *C. philodice* has a similar pattern of winter stage records (Table 1). Williams (377a) states that *cardui* does not survive the U. S. winter in any stage, although many other authors (Table 1) give the adult as the most probable winter stage (as in other *Vanessa*), and Baker (304) thinks it may survive the British winter. *Danaus plexippus* diapauses as an adult, but mainly in Mexico and along the California coast (391, 426); it may not be able to tolerate temperatures far below freezing.

Ecological significance of hibernal diapause stage. The main benefit of a hibernal diapause stage is winter survival. Another significant feature is that the diapause stage is correlated with the seasonal appearance of the adults and other stages during the season. The adult diapause species fly first in spring, followed successively by pupal, mature larval, half-grown larval, young larval, and finally egg diapause species. The first instar diapause species often emerge as late in the season as the egg diapause species, because at least two of the egg diapause species (*H. comma*, *T. lineola*) actually diapause as first-instar larvae inside the egg. This seasonal progression is very striking among the univoltine species and among the first yearly brood of multivoltine species in Colorado (529, 536). It means that insect diapause strategies are adapted to the phenology of their host plants and habitats. The hibernal diapause stage may be adapted to the host plant in other ways. The egg diapause species commonly place their

eggs on the larval host plant where the larvae will not need to search for food in spring. *Lycaena* usually diapause as eggs haphazardly placed around *Rumex* plants. The larval diapause species may or may not need to search for their host plants in spring, depending on where they diapause. The *Limenitis* (*Limenitis*) species diapause as larvae inside a cut-leaf case called a "hibernaculum," usually attached to the tree. *Speyeria* larvae sometimes diapause inside grass stems, and Melitaeini larvae commonly diapause under stones or logs. Many Pyrginae larvae diapause inside their silk tubes constructed from dicotyledon leaves, which fall to the ground in fall. The pupal and adult diapause species do need to search for the host plants in spring. The adult diapause species generally mate in early spring after spending the winter in crevices such as under bark (*Nymphalis*, *Polygonia*) or hanging from trees in dense aggregations (*D. plexippus*). It is conceivable that larval mortality may be high in spring for larvae which must relocate their host plants. The egg diapause species tend to have fewer broods than other species in Colorado; eggs may be better protected against desiccation than other diapause stages except pupae, so may be able to live longer.

Acknowledgements. I thank the persons listed in the literature cited, section D, for providing hibernaculum records. I also thank John F. Emmel, Steven R. Sims, and R. H. T. Mattoni for reviewing the manuscript.

Literature Cited

A. Separate Works (numbers 1-24)

1. BECK, S. D. 1968. Insect photoperiodism. Academic Press, N. Y. and London.
2. CHAPMAN, R. F. 1969. The insects. Structure and function. English Univ. Press, Ltd. London.
3. COMSTOCK, J. A. 1927. Butterflies of California. Publ. by author, Los Angeles, Calif.
4. DÁNILEVSKII, A. S. 1965. Photoperiodism and seasonal development of insects. Oliver & Boyd, Edinburgh.
5. EDWARDS, W. H. 1868-1897. The butterflies of North America. Publ. by author. Philadelphia, Boston, & New York, 3 vol.
6. ESSIG, E. O. 1926. Insects of western North America. MacMillan, New York (p. 635).
7. FORD, E. B. 1962. Butterflies. The New Naturalist Series, Collins, London.
8. FRENCH, G. H. 1899. The butterflies of the eastern United States. Philadelphia.
9. HARRIS, L. H. Jr. Butterflies of Georgia. Univ. Oklahoma Press, Norman, Okla.
10. HOOPER, R. R. 1973. The butterflies of Saskatchewan. Mus. Nat. Hist., Regina, Saskatchewan.
11. HOWARTH, T. G. 1973. South's British butterflies. Frederick Warne & Co., London.
12. HOWE, W., ed., and twenty contributors. 1975. The butterflies of North America. Doubleday, Garden City, N.Y.
13. KLOTS, A. B. 1951. A field guide to the butterflies. Houghton, Mifflin, Cambridge, Mass.

14. LEES, A. D. 1955. The physiology of diapause in arthropods. Cambridge Univ. Press, London, N. Y.
 15. MACY, R. W., & H. H. SHEPARD. 1941. Butterflies. Univ. Minnesota Press, Minneapolis.
 16. MEAD, T. L. 1875. Report upon the collections of diurnal lepidoptera made in portions of Nevada, Utah, Colorado, New Mexico, and Arizona during the years 1871-1874, with notes upon all species known to inhabit Colorado, pp. 739-791, Chapter 8, vol. 5 Zoology, In: G. M. Wheeler & A. A. Humphreys, report upon geological explorations and surveys west of the one hundredth meridian. Washington, D. C.: Government Printing Office.
 17. PYLE, R. M. 1974. Watching Washington butterflies. Seattle Audubon Society, Seattle, Wash.
 18. RAIZENNE, H. 1952. Forest lepidoptera of southern Ontario and their parasites. Canada Dept. Agric. Div. of Forest Biology.
 19. SCUDDER, S. H. 1889. The butterflies of the eastern United States and Canada with special reference to New England. Cambridge, Mass. Pub. by author. 3 vol.
 20. _____. 1893. Brief guide to the commoner butterflies of the northern United States and Canada. Henry Holt & Co., New York.
 21. _____. 1899. Everyday butterflies. Houghton Mifflin Co., Boston.
 22. SHAPIRO, A. M. 1966. Butterflies of the Delaware Valley. Spec. Pub. Amer. Ent. Soc.
 - 22a. TYLER, H. 1975. The swallowtail butterflies of North America. Naturegraph Publ., Healdsburg, California. 192 p.
 23. WEED, C. M. 1923. Butterflies worth knowing. Doubleday, Garden City, N. Y.
 24. WRIGHT, W. G. 1905. The butterflies of the west coast of the United States. Whitaker & Ray Co., San Francisco.
- B. Journals (numbers 25-528)
- Acad. Sci. St. Louis (Missouri). Trans. 25 (H. O'Byrne 1934 28:227).
 Acta Zoologica Fennica. 25a (H. Krogerus 1954 82:3).
 Allyn Museum. Bull. 25 b (R. Wielgus & D. Wielgus 1972 no. 9), 25 c (R. Wielgus & D. Wielgus 1974 no. 17 p. 6), 26 (C. Ferris 1974 no. 18 p. 3), 26 a (R. Wielgus & D. Stallings 1974 no. 23 p. 4), 27 (R. Kendall & W. McGuire 1975 no. 27 p. 3), 28 (R. Kendall 1976 no. 32), 29 (C. Ferris 1977 no. 45 p. 30).
 American Entom. Soc. Trans. 30 (K. Coolidge 1924 50:319), 31 (O. Querci 1936 62:37).
 Amer. Midl. Naturalist. 32 (S. Ae 1958 60:84), 33 (A. Shapiro 1974 91:245), 34 (J. Scott 1974 91:383), 35 (A. Shapiro 1975 93:424), 36 (H. Hill, A. Wenner, P. Wells 1976 95:10).
 Amer. Mus. Nat. Hist. Bull. 37 (W. Beutenmuller 1893 5:241).
 Amer. Mus. Novitates. 38 (H. Freeman 1955 no. 1711 p. 9).
 Amer. Naturalist. 38a (A. Packard 1868 vol. 1), 38b (R. Hoffmann 1978 112:999).
 Ann. Rept. State Entomologist. Missouri. 39 (C. Riley 1876 no. 8 p. 172).
 Ann. Review Ecol. & Systematics. 40 (L. Gilbert 1975 6:384).
 Ann. Review Entom. 40a (M. Tauber & C. Tauber 1976 21:81).
 Assoc. Minnesota Entomol. Bull. 40b (P. Conway 1967 1:43), 41 (W. Krivda 1968 2:48), 42 (J. Masters 1969 3:23).
 Atala. 43 (R. Dirig 1974 2:4), 44 (D. McCorkle 1975 3:9), 45 (R. Pyle 1975 3:34), 46 (L. Orsak 1976 4:13).
 Biol. Medd. Dan. Vid. Selsk. 47 (E. Nielsen 1961 23:1).

(the) Blue Jay. 48 (W. Krivda 1967 25:134).

Brooklyn Entom. Soc. Bull. 49 (C. Rummel 1919 14:47), 50 (K. Coolidge 1924 19:44), 51 (A. Klots 1930 25:147), 52 (R. Leussler & O. Bryant 1935 30:42), 53 (W. McAlpine 1938 33:111), 54 (W. Hovanitz 1941 36:133), 55 (G. Rawson 1945 40:49), 56 (G. Rawson 1951 46:79), 57 (G. Rawson 1951 46:123).

Canada Dept. Agric., Forest Biol. Division. 58 (B. McGugan 1958 Publ. no. 1034).

Canadian Entomologist. 59 (W. Saunders 1868 1:3), 60 (W. Saunders 1868 1:51), 61 (W. Edwards 1868 1:59), 62 (W. Saunders 1868 1:73), 63 (W. Saunders 1868 1:93), 64 (W. Saunders 1869 2:139), 65 (W. Edwards 1869 2:162), 66 (S. Scudder, data of W. Abbot, 1872 4:73, 84), 67 (G. Bowles 1872 4:102), 68 (W. Saunders 1872 4:161), 69 (W. Edwards 1872 4:238), 70 (W. Saunders 1873 5:37), 71 (W. Saunders 1873 5:41), 72 (W. Edwards 1873 5:223), 73 (W. Saunders 1874 6:2), 74 (W. Saunders 1874 6:46), 75 (W. Edwards 1874 6:121), 76 (W. Edwards 1874 6:159), 77 (W. Edwards 1875 7:80), 78 (F. Caulfield 1875 7:86), 79 (W. Edwards 1875 7:150), 80 (W. Edwards 1875 7:189), 81 (W. Edwards 1875 7:228), 82 (S. Scudder 1876 8:61), 83 (W. Edwards 1876 8:119), 84 (W. Edwards 1876 8:148), 85 (W. Edwards 1877 9:1), 86 (W. Edwards 1877 9:34), 87 (F. Caulfield 1877 9:40), 88 (W. Edwards 1877 9:114), 89 (W. Edwards 1877 9:141), 90 (W. Edwards 1877 9:165), 91 (W. Edwards 1878 10:1), 92 (W. Edwards 1878 10:69), 93 (W. Edwards 1878 10:105), 94 (C. Siewers 1878 10:115), 95 (J. Doll 1878 10:154), 96 (W. Edwards 1879 11:31), 97 (W. Edwards 1879 11:101), 98 (W. Edwards 1879 11:127), 99 (W. Edwards 1879 11:141), 100 (W. Edwards 1879 11:177), 101 (J. Morris 1879 11:200), 102 (W. Edwards 1879 11:217), 103 (W. Edwards 1879 11:224), 104 (C. Heustis 1879 11:239), 105 (W. Edwards 1880 12:9), 106 (W. Edwards 1880 12:21), 107 (W. Edwards 1880 12:69), 108 (W. Murray 1880 12:120), 109 (W. Edwards 1880 12:141), 110 (W. Edwards 1880 12:231), 111 (W. Edwards 1881 13:61), 112 (W. Edwards 1881 13:81), 113 (J. Moffat 1881 13:115), 114 (W. Edwards 1881 13:119), 115 (W. Edwards 1881 13:177), 116 (W. Edwards 1881 13:205), 117 (W. Edwards 1881 13:226), 118 (W. Edwards 1881 13:237), 119 (W. Saunders 1882 14:1), 120 (W. Edwards 1882 14:20), 121 (W. Edwards 1882 14:29), 122 (W. Edwards 1882 14:49), 123 (W. Edwards 1882 14:84), 124 (E. Reed 1882 14:160), 125 (W. Edwards 1882 14:189), 126 (W. Edwards 1882 14:201), 127 (W. Edwards 1882 14:229), 128 (A. Grote 1883 15:40), 129 (P. Gosse 1883 15:44), 130 (W. Edwards 1883 15:64), 131 (W. Edwards 1884 16:81), 132 (W. Edwards 1884 16:109), 133 (W. Edwards 1884 16:163), 134 (W. Edwards 1884 16:232), 135 (W. Edwards 1885 17:61), 136 (W. Edwards 1885 17:98), 137 (W. Edwards 1885 17:108), 138 (W. Edwards 1885 17:155), 139 (W. Edwards 1885 17:245), 140 (W. Edwards 1886 18:14), 141 (W. Edwards 1886 18:88), 142 (W. Edwards 1886 18:141), 143 (W. Edwards 1886 18:201), 144 (W. Edwards 1887 19:34), 145 (W. Edwards 1887 19:41), 146 (S. Scudder 1887 19:101), 147 (W. Edwards 1887 19:170), 148 (W. Edwards 1887 19:226), 149 (W. Edwards 1888 20:1), 150 (W. Edwards 1888 20:3), 151 (W. Edwards 1888 20:41), 152 (W. Marsh 1888 20:45), 153 (W. Edwards 1888 20:67), 154 (W. Edwards 1888 20:83), 155 (T. Fyles 1888 20:131), 156 (C. Bethune 1888 20:216), 157 (E. Morton 1888 20:226), 158 (T. Fyles 1889 21:12), 159 (W. Edwards 1889 21:21), 160 (W. Edwards 1890 22:62), 160a (D. Bruce 1890 22:67), 161 (A. Winn 1890 22:80), 162 (T. Bean 1890 22:94), 163 (W. Patton 1891 23:66), 164 (H. Dyar 1891 23:172 + 201), 165 (A. Bethune 1891 23:200), 166 (H. Lyman 1892 24:1), 167 (W. Edwards 1892 24:49), 168 (J. Fletcher 1893 25:87), 169 (R. Wolcott 1893 25:98), 170 (W. Edwards 1893 25:253), 171 (J. Lambert 1893 25:259), 172 (W. Edwards 1893 25:286), 173 (W. Edwards 1894

26:3), 174 (W. Harrington 1894 26:9), 175 (J. Fletcher 1894 26:27), 176 (W. Edwards 1894 26:55), 177 (E. Heath 1895 27:310), 178 (H. Lyman 1895 27:333), 179 (T. Fyles 1895 27:346), 180 (A. Snyder 1896 28:105), 181 (H. Lyman 1896 28:143), 182 (W. Fiske 1896 28:162), 183 (W. Fiske 1897 29:26), 184 (A. Snyder 1897 29:118), 185 (H. Lyman 1897 29:249), 186 (H. Lyman & A. Winn 1897 29:273), 187 (A. Good 1901 33:228), 188 (H. Lyman 1902 34:126), 189 (J. Cook & H. Cook 1904 36:136), 190 (J. Fletcher 1905 37:79), 191 (T. Fyles 1906 38:137), 192 (J. Cook 1906 38:141), 193 (J. Cook 1906 38:181), 194 (J. Cook 1906 38:214), 195 (J. Cook 1907 39:145), 196 (J. Cook 1907 39:229), 197 (J. Cook 1907 39:293), 198 (J. Cook 1907 39:405), 199 (J. Cook & F. Watson 1908 40:85), 200 (F. Wolley-Dod 1908 40:149), 201 (F. Wolley-Dod 1908 40:181), 202 (K. Coolidge 1908 40:425), 203 (W. Newcomb 1909 41:221), 204 (A. Gibson 1910 42:145), 205 (E. Newcomer 1911 43:83), 206 (W. Newcomb 1911 43:160), 207 (A. Winn 1912 44:213), 208 (T. Cockerell 1915 47:280), 209 (J. McDunnough 1920 52:56), 210 (J. McDunnough 1922 54:134), 211 (H. Mousley 1923 55:26), 212 (J. McDunnough 1930 62:107), 213 (R. Handford 1933 65:95), 214 (C. dos Passos & L. Grey 1934 66:188 & 278), 215 (C. dos Passos 1935 67:82), 216 (C. dos Passos 1936 68:167), 217 (C. dos Passos 1936 68:239), 218 (V. Dethier 1938 70:255), 219 (V. Dethier 1939 71:117), 220 (J. McDunnough 1939 71:153), 221 (J. McDunnough 1940 72:130), 222 (G. Wood 1951 83:241), 223 (V. Dethier 1959 91:581), 223a (F. Brown 1961 93:107), 223b (J. Burns 1966 98:859).

Carnegie Museum. Ann. 224 (J. Bonniwell 1931 20:264).

Cornell Univ. Agric. Exp. Stn., Ithaca, N.Y. Memoirs. 225 (W. Forbes 1960 no. 371).

Denver (Colorado) Museum Nat. Hist. Proc. 226 (F. Brown, D. Eff, B. Rotger 1957 no. 3-7).

Ecol. Entom. 227 (J. Dempster, M. King, K. Lakhani 1976 1:71), 228 (A. Shapiro 1977 2:217), 228a (P. Tuskes & L. Brower 1978 3:141).

Ecology. 229 (P. Brussard & P. Ehrlich 1970 51:119).

Entom. Scandinavica. 230 (J. Scott 1973 4:161).

Entom. Monthly Mag. 231 (B. Petersen 1962 98:200).

Entom. News. 232 (W. Osborne 1895 6:245 & 281), 233 (D. Bruce 1896 7:15), 234 (E. Smyth Jr. 1900 11:465), 235 (W. Newcomb 1900 11:514), 236 (W. Field 1900 11:514), 237 (J. Cook 1903 14:143), 238 (C. Brimley & F. Sherman Jr. 1903 14:230), 239 (J. Grundel 1904 15:97), 240 (J. Grundel 1905 16:86), 241 (J. Grossbeck 1905 16:131), 242 (P. Laurent 1905 16:245), 243 (P. Laurent 1906 17:103), 244 (R. Rowley 1906 17:175), 245 (C. Shull 1907 18:73), 246 (R. Rowley 1907 18:306), 247 (F. Wolley-Dod 1908 19:114), 248 (E. Smyth Jr. 1908 19:191), 249 (P. Laurent 1908 19:408), 250 (F. Williams 1908 19:476), 251 (F. Williams 1909 20:62), 252 (W. Newcomb 1909 20:293), 253 (F. Williams 1910 21:30), 254 (F. Williams 1910 21:279), 255 (H. Bower 1911 22:359), 256 (P. Laurent 1912 23:483), 257 (G. Bethune-Baker 1913 24:97), 258 (G. Bethune-Baker 1913 24:152), 259 (A. Girault 1913 24:195), 260 (R. Leussler 1913 24:344), 260a (P. Schroers 1913 24:355), 261 (H. Skinner 1914 25:326), 262 (J. Franzen 1914 25:363), 263 (J. Bonniwell 1916 27:372), 264 (K. Coolidge 1922 33:305), 265 (K. Coolidge 1923 34:295), 266 (K. Coolidge 1924 35:115), 267 (K. Coolidge 1924 35:306), 268 (K. Coolidge 1924 35:199), 269 (K. Coolidge 1925 36:65), 270 (S. Bromley 1928 39:96), 271 (R. Leussler 1930 41:7), 272 (V. Calkins 1932 43:210, 225, 257), 273 (V. Kite 1934 45:36), 274 (T. Cockerell 1934 45:174), 275 (E. Meiners 1936 47:269), 276 (R. Leussler 1938 49:3, 76, 213, 275, 50:34), 277 (H. O'Byrne 1941 52:181), 278 (C. Wood Jr. & C. Gottschalk 1942 53:143, 159, 191), 279 (F. Brown 1952 63:122), 280

(P. Ehrlich & H. Clench 1960 71:137), 281 (J. Downey 1962 73:57), 282 (J. Masters 1978 79:85), 283 (A. Shapiro 1969 80:130), 284 (R. Anderson 1971 82:5), 285 (A. Shapiro 1971 82:13), 286 (W. Krivda 1972 83:103), 286a (C. Turnbull 1979 90:125).

Entom. Soc. Amer. Ann. 287 (A. Holmquist 1926 19:395), 288 (V. Randolph 1927 20:242), 289 (R. Barker, A. Mayer, C. Cohen 1963 56:292), 290 (I. Nakamura & S. Ae 1977 70:481).

Entom. Soc. Ontario. Ann. Report. 291 (J. Fletcher 1888 no. 19 p. 74).

Entom. Soc. Ontario. Proc. 292 (D. Pengelly 1961 91:189).

(the) Entomologist. 293 (S. Bowden 1966 99:175).

Evolution. 294 (S. Ae 1959 13:65), 295 (A. Shapiro & R. Carde 1970 24:53), 296 (M. Singer 1971 25:383), 296a (J. Brittnacher, S. Sims, F. Ayala 1978 32:199), 296b (M. Douglas & J. Grula 1978 32:776), 296c (C. Oliver 1972 26:221).

Florida Dept. Agric. Arthropods of Florida. 297 (C. Kimball 1965 v. 1).

Great Basin Naturalist. 297a (A. Shapiro 1977 37:443).

Hilgardia. 298 (V. Stern & R. Smith 1960 29:411).

Indiana State Geologist. Annual Report. 299 (W. Blatchley 1891 17:365).

Insect Biochemistry. 299a (R. Cenedella 1971 1:244).

Int. Congr. Entom. no. 10. Proc. 300 (C. Remington 1958 2:787).

J. Agric. Univ. Puerto Rico. 301 (G. Wolcott 1943 27:93), 302 (G. Wolcott 1948 32:561).

J. Anim. Ecol. 303 (R. Jones 1977 46:195).

J. Comp. Physiol. 303a (W. Herman & D. Bennett 1975 99:331).

J. Entomol. (A). 304 (R. Baker 1972 46:185).

J. Insect Physiol. 305 (C. Oliver 1969 15:1579), 306 (S. Clark & A. Platt 1969 15:1951), 306a (W. Herman 1973 19:1883), 306b (J. Brower & G. Chippendale 1974 20:1117), 306c (J. Hong & A. Platt 1975 21:1159).

J. Research Lepid. 307 (J. Garth & J. Tilden 1963 2:1), 308 (F. Thorne 1963 2:146), 309 (J. Heitzman 1963 2:170), 310 (E. Newcomer 1963 2:279), 310b (G. N. Ross 1964 3:9), 310a (E. Newcomer 1964 3:49), 311 (K. Roever 1964 3:103), 312 (F. Brown 1964 3:121), 313 (H. Freeman 1964 3:145), 314 (J. Heitzman 1964 3:151), 315 (J. Heitzman 1964 3:154), 315a (L. LaDue 1964 3:197), 316 (J. Heitzman 1965 4:75), 317 (E. Newcomer 1966 5:243), 318 (A. Shapiro 1967 6:181), 318a (J. Masters 1968 7:19), 319 (T. Emmel & J. Emmel 1968 7:125), 320 (T. Emmel 1968 7:159), 321 (J. Heitzman & R. Heitzman 1969 8:193), 322 (R. Simmons & W. Andersen 1970 9:181), 323 (J. Emmel, O. Shields, R. Breedlove 1970 9:238), 324 (C. Callaghan & K. Tidwell 1972 10:198), 325 (W. McAlpine 1972 11:83), 326 (R. Wielgus & D. Wielgus 1972 11:249), 327 (A. Shapiro & A. Shapiro 1973 12:65), 327a (J. Scott 1973 12:151), 328 (J. Scott 1973 12:225), 328a (A. Shapiro 1974 13:73), 329 (J. Heitzman & R. Heitzman 1974 13:241), 329a (C. Ferris 1974 13:249), 330 (R. Neck 1975 14:42), 330a (C. Ferris 1976 15:1), 331 (J. Tilden 1976 15:33), 331a (C. Ferris 1976 15:106), 331b (A. Shapiro 1977 16:83), 332 (J. Tilden & D. Huntzinger 1977 16:176).

Kansas Entom. Soc. Journal. 332a (J. Masters 1969 42:133), 333 (C. Goodpasture 1973 46:477).

Kontyu (Entomology). 334 (T. Sakai & S. Masaki 1965 33:275).

(the) Lepidopterist's Society. Journal. 335 (R. Guppy 1948 2:supplement pp. iii), 335a (V. Hynes 1949 3:64), 336 (F. Thorne 1949 3:86), 337 (J. Hopfinger 1949 3:88), 338 (W. Evans 1949 3:106), 338a (W. Evans 1950 4:70), 338b (D. Bauer 1950 4:90), 339 (D. Eff 1950 4:94), 340 (W. Howe 1950 4:96), 41 (H. Chase 1950 4:98), 342 (D. Downey 1951 5:92), 342a (H. Chase 1951 5:99), 343 (A. Brower 1951

5:107), 343a (W. Evans 1952 6:106), 344 (J. Tilden 1953 7:69), 344a (R. Guppy 1953 7:85), 345 (P. Remington 1953 7:98), 346 (J. Merritt 1953 7:101), 347 (W. Randle 1953 7:119), 348 (T. Iwase 1954 8:95), 348a (V. Muspratt 1954 8:143), 348b (R. Guppy 1955 9:15), 348c (W. Reinthal 1956 10:25), 348d (E. Meiners 1956 10:163), 349 (G. Ehle 1957 11:39), 349a (C. Wyatt 1957 11:46), 350 (D. Stallings & J. Turner 1957 11:114), 350a (J. Tilden 1957 11:230), 350b (J. Tilden 1958 12:32), 351 (M. Nielsen 1958 12:37), 351a (A. Brower 1958 12:47), 352 (G. Hardy 1959 13:70), 353 (C. Remington 1959 13:164), 354 (W. McAlpine, S. Hubbell, T. Pliske 1960 14:209), 355 (J. Downey & W. Fuller 1961 15:40), 356 (J. Lipes 1961 15:114), 357 (H. Clench 1961 15:226), 357a (Z. Lorkovic 1962 16:5), 358 (T. Emmel & J. Emmel 1962 16:31), 359 (D. Stallings, J. Turner, P. Ehrlich 1962 16:56), 360 (F. Thorne 1962 16:61), 361 (W. Reinthal & R. Kendall 1962 16:202), 362 (R. Kendall 1963 17:22), 363 (D. Chambers 1963 17:26), 363a (R. Heitzman 1963 17:44), 364 (G. Ross & E. Lambremont 1963 17:157), 364a (R. Turner 1963 17:227), 365 (S. Bowden 1964 18:95), 366 (R. Kendall 1964 18:129), 367 (R. Kendall 1964 18:163), 368 (R. Guppy 1964 18:187), 369 (E. Newcomer 1964 18:217), 370 (R. Kendall 1965 19:1), 370a (W. Howe 1965 19:33), 370b (A. Legge 1965 19:57), 371 (J. Heitzman 1965 19:79), 372 (R. Kendall 1966 20:35), 373 (R. Kendall 1966 20:229), 374 (E. Newcomer 1967 21:71), 374a (J. Masters 1967 21:258), 374b (J. Emmel & T. Emmel 1968 22:46), 374c (W. Krivda 1968 22:191), 375 (R. Langston 1969 23:149), 375a (J. Masters & J. Sorensen 1969 23:155), 376 (T. Emmel 1969 23:172), 376a (J. Muller 1969 23:249), 377 (C. Oliver 1970 24:78), 377a (C. Williams 1970 24:157), 377b (F. Rutkowski 1971 25:126), 377c (J. Masters 1971 25:150), 377d (B. Drummond, G. Bush, T. Emmel 1970 24:136), 378 (S. Mattoon, R. Davis, O. Spencer 1971 25:248), 379 (G. Gorelick 1971 25:supplement 2), 379a (E. Shull & F. Badger 1972 26:13), 380 (T. Emmel, S. Mattoon 1972 26:143), 380a (P. Kean & A. Platt 1973 27:112), 381 (A. Shapiro 1973 27:158), 382 (C. Ferris 1973 27:279), 383 (C. Goodpasture 1974 28:55), 384 (S. Ellis 1974 28:120), 384a (A. Muyshondt 1974 28:174), 385 (J. Masters 1974 28:237), 386 (J. Emmel & T. Emmel 1974 28:346), 387 (A. Shapiro 1974 28:372), 387a (W. Evans 1975 29:53), 388 (J. Scott 1975 29:166), 389 (R. Kendall & M. Rickard 1976 30:105), 389a (N. Proctor 1976 30:126), 390 (P. Ehrlich 1976 30:149), 391 (F. Urquhart & N. Urquhart 1976 30:153), 391a (A. Shapiro 1976 30:289), 392 (W. Wagner Jr. & T. Mellichamp 1978 32:29), 393 (T. Dimock 1978 32:94), 393a (M. Bowers 1978 32:282), 393b (A. Platt, G. Rawson, G. Balogh 1978 32:289), 393c (C. Oliver 1979 33:6), 393d (J. Masters 1979 33:167).

(the Lepidopterist's Society. News. 394 (R. Langston 1961 no. 4 p. 2), 395 (R. Langston & J. Tilden 1962 no. 3 p. 2), 395a (R. Guppy 1962 no. 3 p. 3), 396 (W. Reinthal 1962 no. 3 p. 10), 397 (P. Herlan 1963 no. 4 p. 3), 398 (B. Perkins & S. Perkins 1964 no. 4 p. 4), 398a (D. Eff 1964 no. 4 p. 5), 398b (H. Hensel 1965 no. 3 p. 10), 398c (J. Legge 1966 no. 3 p. 6), 399 (R. Heitzman 1966 no. 3 p. 9), 400 (L. Paulissen 1966 no. 3 p. 10), 401 (M. Nielsen 1967 no. 3 p. 11), 402 (J. Heitzman 1968 no. 3 p. 11), 403 (J. Mori 1969 no. 3 p. 7), 404 (J. Masters 1969 no. 3 p. 13), 405 (K. Philip & L. Miller 1969 no. 3 p. 20), 406 (K. jPhilip 1970 no. 3 p. 18), 407 (K. Philip 1971 no. 3 p. 6), 408 (K. Philip & C. Ferris 1972 no. 2 p. 15), 408a (D. Schweitzer 1973 no. 2 p. 16), 409 (K. Philip 1973 no. 2 p. 18), 410 (Anonymous 1970 no. 3 p. 13), 411 (K. Hansen 1972 no. 2 p. 2), 412 (B. Griffin 1973 no. 3 p. 7), 413 (R. Dirig 1974 no. 2 p. 13), 413a (L. Hoying 1975 no. 2/3 p. 9), 413b (R. Boscoe 1975 no. 2/3 p. 15), 414 (D. Eff & R. Pyle 1975 no. 2/3 p. 6), 414a (P. Grey 1975 no. 2/3 p. 13), 414b (C. Bird 1976 no. 2 p. 5), 414c (C. Ferris 1976 no. 3 p. 1), 415 (Anonymous 1977 no. 1 p. 6), 416 (A. Shapiro 1977 no. 2 p. 5), 417 (L. Paulissen 1977 no. 2 p. 13),

417a (R. Layberry 1977 no. 2 p. 15), 418 (T. Williams 1977 no. 2 p. 18), 419 (D. Winter 1977 no. 2 p. 19), 419a (K. Philip 1977 no. 4 p. 4), 420 (R. Langston 1978 no. 2 p. 4), 421 (C. Oliver 1978 no. 2 p. 4), 422 (R. Boscoe 1978 no. 2 p. 14), 422a (Q. Hess 1978 no. 2 p. 16), 422d (R. Langston 1979 no. 2 p. 4), 422e (M. Smith, S. Mattoon 1979 no. 2 p. 5), 422f (F. Lee 1979 no. 2 p. 9).

Mid-Continent Lepidoptera Series. 422b (J. Masters 1971 no. 25), 422c (J. Masters 1972 no. 52).

Milwaukee (Wisconsin) Public Museum. Popular Science Handbooks. 423 (J. Ebner 1970 no. 12).

Museum Comp. Zool. Bull. 424 (D. Davenport 1941 87:263).

National Geographic. 425 (F. Urquhart 1976 150:161).

Natural History. 426 (L. Brower 1977 86:41).

Natural Hist. Mus. Los Angeles County. Science Series. 427 (T. Emmel & J. Emmel 1973 26:1).

Nature. 428 (V. Dethier & R. MacArthur 1964 201:728).

New York Acad. Sci. Survey Puerto Rico. 429 (W. Comstock 1944 12:481).

New York Entom. Soc. Journal. 430 (W. Davis 1893 1:43), 431 (H. Skinner 1896 4:107), 432 (A. Klots 1935 43:139), 433 (W. Comstock 1940 48:47), 434 (F. Brown 1958 66:63), 435 (A. Klots 1971 79:84), 436 (A. Shapiro 1971 79:27).

New York State Museum (Albany). Handbooks. 437 (A. Saunders 1932 no. 13:1).

Nova Scotia Museum Science Bull. 438 (D. Ferguson 1955 2:161).

Oecologia. 439 (R. White 1974 14:309), 439a (W. Watt, D. Han, B. Tabashnik 1979 44:44).

Ottawa Naturalist. 440 (J. Williams 1902 January).

Pan-Pacific Entom. 441 (R. Langston & J. Comstock 1966 42:104), 442 (J. Scott 1974 50:20).

Papilio. 443 (Anonymous 1881 1:125), 444 (W. Edwards 1881 1:139), 445 (W. Edwards 1881 1:151), 446 (J. Rivers 1883 3:26), 447 (W. Edwards 1883 3:90), 448 (A. Grote 1883 3:123), 449 (W. Edwards 1883 3:159), 450 (W. Edwards 1884 4:63), 451 (W. Wright 1884 4:126), 452 (W. Edwards 1884 4:131), 453 (W. Edwards 1884 4:166).

Physiology of Insecta. M. Rockstein, ed. 454 (M. Keister & J. Buck 1974 6:469).

Psyche. 455 (S. Scudder 1874 1:13), 456 (S. Scudder 1874 1:18), 457 (W. Edwards 1880 3:3), 458 (W. Edwards 1880 3:75), 459 (W. Edwards 1881 3:159, 171), 460 (W. Edwards 1881 3:174), 461 (W. Dwight 1882 3:327), 462 (W. Edwards 1886 4:353), 463 (H. Hinkley 1888 5:54), 464 (S. Scudder 1890 5:330), 465 (S. Scudder 1888 5:63), 466 (S. Scudder 1889 5:137), 467 (H. Weed 1894 7:129), 468 (N. Durand 1900 9:87), 469 (S. Scudder 1901 9:195), 470 (K. Coolidge 1907 14:118), 471 (K. Coolidge 1908 15:81), 472 (W. Field 1914 21:115), 473 (C. Remington 1952 59:61), 474 (R. Carde, A. Shapiro, H. Clench 1970 77:70), 475 (J. Burns 1974 81:51), 476 (F. Slansky 1974 81:243), 477 (J. Kanz 1977 84:120), 477a (A. Shapiro 1979 86:1).

San Diego Soc. Nat. Hist. Trans. 478 (J. Comstock & R. Kendall 1967 14:233).

Science. 479 (M. Singer 1972 176:75), 480 (P. Ehrlich, R. White, M. Singer, S. McKechnie, L. Gilbert 1975 188:221).

Search. (Cornell Univ. Agric. Exp. Stn.) Entomology. 481 (A. Shapiro 1974 12:1).

Smithsonian Inst. Annual Report. 481a (A. Clark 1925, Publ. 2856 p. 439).

Smithsonian Misc. Collns. Wash. D. C. 482 (A. Clark & L. Clark 1951 116:1).

Southern Calif. Acad. Sci. Bull. 483 (J. Comstock 1928 27:64), 484 (J. Comstock 1929 28:52), 485 (J. Comstock 1930 29:141), 486 (J. Comstock & C. Dammers 1931 30:22), 487 (J. Comstock & C. Dammers 1932 31:13), 488 (J. Comstock & C. Dammers 1932 31:33), 489 (J. Comstock & C. Dammers 1932 31:88), 490 (J. Comstock & C. Dammers 1933 32:28), 491 (J. Comstock, G. Sperry, & J. Sperry 1933 32:100), 492 (J. Comstock & C. Dammers 1933 32:105), 493 (J. Comstock 1933 32:114), 494 (J. Comstock & C. Dammers 1934 33:25), 495 (J. Comstock & C. Dammers 1934 33:79), 496 (J. Comstock & C. Dammers 1934 33:137), 497 (J. Comstock & C. Dammers 1935 34:82), 498 (J. Comstock & C. Dammers 1935 34:120), 499 (J. Comstock 1935 34:148), 500 (J. Comstock & C. Dammers 1935 34:211), 501 (J. Comstock & C. Dammers 1936 35:8), 502 (J. Comstock, J. Sperry, & G. Sperry 1936 35:165), 503 (J. Comstock & C. Dammers 1937 36:18), 504 (J. Comstock & C. Dammers 1938 37:32), 505 (V. Dethier 1938 37:76), 506 (V. Dethier 1939 38:190), 507 (V. Dethier 1940 39:110), 508 (J. Comstock 1940 39:113), 509 (C. Dammers 1940 39:124), 510 (V. Dethier 1941 40:109), 511 (V. Dethier 1942 41:41), 512 (V. Dethier 1944 43:30), 513 (J. Comstock 1955 54:59), 514 (J. Downey & W. Lange 1956 55:153), 515 (J. Comstock 1956 55:178), 516 (J. Comstock 1961 60:147).

St. Louis Acad. Sci. Trans. 517 (C. Riley 1877 3:567, 323).

Texas Agric. Exp. Stn. Bull. 518 (H. Reinhard 1929 no. 401 p. 32).

U. S. Nat. Mus. Bull. 519 (A. Clark 1932 157:1).

Univ. Calif. Press. Berkeley Calif. Calif. Nat. Hist. Guides. 520 (J. Tilden 1965 no. 12:1).

Univ. Calif. Publ. Entom. 521 (C. MacNeill 1964 35:1), 522 (J. Burns 1964 37:1).

Univ. Maine. Life Sciences & Exp. Stn. at Orono. 523 (A. Brower 1974 Tech. Bull. no. 66).

Univ. Montana. Bull. no. 30, Biol. Series no. 10. 524 (M. Elrod 1906).

Wasmann J. Biol. 525 (W. Hovanitz 1951 9:1), 526 (A. Shapiro 1968 26:137), 527 (A. Shapiro 1973 31:291), 528 (A. Shapiro 1974 32:267).

C. Personal records of J. Scott. Number 529. Most records are for Colorado; some are for California and elsewhere.

D. Records based on personal communication (numbers 530-554). 530 Richard A. Arnold, 531 Auburn E. Brower, 532 Lincoln P. Brower, 533 F. Martin Brown, 533a Robert P. Dana, 534 Scott L. Ellis, 535 John F. Emmel, 536 Marc Epstein, 537 Clifford D. Ferris, 538 Michael S. Fisher, 539 Jack L. Harry, 540 J. Richard Heitzman, 541 Roy O. Kendall, 542 Sterling O. Mattoon, 543 Lee D. Miller, 543a Douglas Mullins (pers. comm. to John F. Emmel), 544 Erval J. Newcomer, 545 Kilian Roeber, 546 Arthur M. Shapiro, 547 Steven R. Sims, 548 Michael C. Singer, 549 Ray E. Stanford, 550 Fred T. Thorne, 551 David L. Wagner, 552 Ralph E. Wells, 553 Michael Young, 554 Steven Stone.

Table 1

Hibernal diapause stage for temperate zone species which probably have a true diapause. Diapause stage: E-egg, L-larva (instar given in arabic numbers, or y-young larva, h-part or about half grown larva, m-mature larva), P-pupa, A-adult. Diapause stage for biennial species is indicated by I-first winter, and II-second winter. e-probably an error; diapause probably occurs in the other stage listed for the species (some sources cited after an e may prove that diapause does not occur in this stage). r-winter rarely spent in this stage.

SPECIES	HIBERNAL DIAPAUSE STAGE	SOURCE
Papilionidae		
<i>Parnassius phoebus</i>	E-I	473, 449, 230, 5, 99, 131, 135, 536, 16, 529
	L (h, nearly m)-II	5, 536, 330a, 160a (L or P), 22a
<i>Parnassius clodius</i>	E-I	449, 99, 131, 135
	L (h or m)-II	535, 22a
<i>Parnassius eversmanni</i>	E-I	348
	P-II	348
<i>Papilio glaucus</i>	P	13, 9, 19, 20, 21, 540, 433, 5, 23, 58, 234, 431, 232, 234, 73, 62, 66
<i>Papilio glaucus rutulus</i>	P	532, 58, 546, 5, 15, 167
<i>Papilio eurymedon</i>	P	532
<i>Papilio multicaudata</i>	P	532, 272, 366, 5, 167, 251, 554
<i>Papilio indra</i>	P	427, 412, 310a, 374b
<i>Papilio bairdii</i>	P	5, 549, 170, 310a
<i>Papilio polyxenes</i>	P	13, 366, 20, 21, 23, 237, 540, 101, 220, 234, 275, 305, 15, 276, 433, 19, 8, 232, 234
	A(e)	21, 19
<i>Papilio polyxenes rudkini</i>	P	427, 503, 477a
<i>Papilio polyxenes kahli</i>	P	300
<i>Papilio (polyxenes?) joanae</i>	P	540
<i>Papilio machaon</i>	P	7, 52, 13, 227, 305
<i>Papilio brevicauda</i>	P	13, 23, 19, 129, 220
<i>Papilio androgeus</i>	P	310b
<i>Papilio zelicaon</i>	P	5, 538, 453, 305, 315a, 335a
<i>Papilio cresphontes</i>	P	19, 20, 21, 461, 23, 433, 540, 115, 113, 108, 232, 95, 104
<i>Papilio aristodemus</i>	P	13, 377b
<i>Papilio xuthus</i>	P	390
<i>Papilio troilus</i>	P	13, 366, 9, 540, 433, 23, 468, 19, 20, 21, 232, 66

<i>Eurytides marcellus</i>	P	13, 366, 19, 20, 21, 458, 252, 248, 246, 540, 81, 120, 5, 234, 8, 244, 433, 23, 457, 66, 234, 232, 65, 342a
<i>Battus philenor</i>	P	19, 433, 547, 13, 540, 519, 21, 23, 232, 276
	A(e?)	19, 13, 20, 21, 23
<i>Battus polydamas</i>	P	366, 384a
Pieridae		
<i>Neophasia menapia</i>	E	6, 427, 5, 15, 24, 58, 529
<i>Anthocaris sara</i>	P	427, 520, 471, 5, 167, 343a, 387a, 477a
<i>Anthocaris midea</i>	P	13, 19, 20, 540, 5, 15, 433, 23, 234, 144, 167, 241, 541
<i>Anthocaris lanceolata</i>	P	5, 421, 251, 254, 477a
<i>Anthocaris cethura</i>	P	167, 488, 477a
<i>Anthocaris cethura pima</i>	P	471
<i>Euchloe ausonides</i>	P	5, 529, 167, 327a, 477a
<i>Euchloe olympia</i>	P	13, 12, 540, 15, 245
<i>Euchloe hyantis</i>	P	427, 269, 488, 477a
<i>Pieris napi</i>	P	357a, 228, 13, 520, 3, 70, 129, 19, 20, 23, 431, 231, 365, 14, 4, 1, 377
<i>Pieris rapae</i>	P	191, 13, 433, 482, 289, 318, 540, 348, 23, 8, 15, 431, 374c, 331b, 19, 20, 21, 9, 67, 71, 303, 35, 289, 231, 232, 1, 328a
<i>Pieris occidentalis</i>	P	35, 527, 391a
<i>Pieris protodice</i>	P	13, 299, 318, 20, 19, 35, 23, 8, 21, 540, 433, 119, 71, 55, 31, 526, 232, 546, 391a
<i>Pieris virginiensis</i>	P	285, 13, 433, 12, 432, 413, 293, 348c
<i>Pieris sisymbrii</i>	P	477a, 535
<i>Eurema nicippe</i>	A	21, 20, 540, 519, 12, 299, 22, 278, 111, 19
	P?	232
<i>Colias meadii</i>	L(1)-I	162
	L(3-4, 5[r])-II	12, 226, 32, 5, 167, 147, 159
<i>Colias scudderii</i>	L(3-4)	226, 32, 5
<i>Colias scudderii gigantea</i>	L(y, 3)	12, 210
<i>Colias palaeno</i>	L(3)	12
<i>Colias alexandra</i>	L(2-4)	12, 226, 32, 5, 462, 384, 137, 147, 148
<i>Colias (alexandra?) harfordii</i>	L	5, 147
<i>Colias interior</i>	L(1, 2, 3)	13, 12, 222, 32, 185, 422f
<i>Colias nastes</i>	L(y, 3, m)- (I+II?)	13, 12

<i>Colias eurytheme</i>	L(3-4)	328a, 298, 482, 147, 433, 327, 318, 23, 21, 5, 546, 15, 19, 20
	P	482, 13, 299, 318, 5, 15, 232
	A(e)	482, 13, 299, 21, 15, 19, 20
<i>Colias philodice</i>	L(3-4)	19, 225, 13, 147, 166, 299, 433, 20, 525, 449, 21, 294, 32, 5, 15, 19, 439a, 38b
	P	13, 299, 48, 232
	A(e)	13
<i>Colias hecla</i>	L-I+II	12, 13
	P?	13
Libytheidae		
<i>Libytheana bachmanii</i>	P	13, 225, 23
	A	15, 117, 19, 433, 232
Nymphalidae		
<i>Coenonympha tullia</i>	L(3-4, 5[r])	12, 434, 312, 424, 167, 140, 143, 145, 338a, 223a
<i>C. tullia californica</i>	L(1)	338, 40
<i>C. tullia heinemani</i>	L(1?, 2, h?, m?)	434, 223a
<i>Cercyonis pegala</i>	L(1)	13, 299, 12, 21, 23, 5, 89, 106, 20, 132, 8, 536, 19, 15, 433, 540, 284, 423, 459, 516, 376, 380
<i>Cercyonis meadii</i>	L(1)	226, 12, 5, 376
<i>Cercyonis oetus</i>	L(1)	137, 376, 141, 226, 462, 12, 5
<i>Cercyonis stheneke</i>	L(1)	376, 12, 338a
<i>Erebia magdalena</i>	L(h)	529
<i>Erebia theano</i>	L(y)+?-I&II	529
<i>Erebia epipsodea</i>	L(3-4)	12, 226, 229, 5, 529
<i>Erebia discoidalis</i>	L(4)	5
	P(e)	41, 286
<i>Euptychia areolata</i>	L(h)	225, 13, 5, 433, 19
<i>Euptychia areolata mitchelli</i>	L(4)	354, 12
<i>Euptychia rubricata</i>	L	515
<i>Euptychia cymela</i>	L(2[r], 4, m[r])	13, 299, 366, 423, 433, 225, 64, 20, 284, 21, 93, 23, 15, 8, 540, 19, 93, 429, 232
<i>Cyllopsis gemma</i>	L(4)	225, 366, 540, 5, 19, 232, 96, 130
<i>Cyllopsis pertepida</i>	L(h)	529
<i>Oeneis alberta</i>	L(m)?	502, 5
	P?	279
<i>Oeneis uhleri</i>	L(2, 3, 4, 5)	5, 12, 226
<i>Oeneis melissa</i>	L(y, 2, 3)-I	5, 469, 23, 225, 15, 19
	L(5)-II	5, 23, 225, 19
<i>Oeneis polixenes</i>	L(1, perhaps 2-3)-I	226, 12, 5
	L(4, 5)-II	5

<i>Oeneis jutta</i>	L(1, 2, 3)-I	19, 158, 12, 226, 225, 406, 5, 168
	L(4, 5, 6)-II	19, 155, 15, 225, 5, 168
<i>Oeneis nevadensis</i>	L(2, 3)-I	5, 176
	L(5)-II	5
<i>Oeneis macounii</i>	L(1, 2)-I	225, 5, 19
	L(m)-II	225, 5, 168
<i>Oeneis chryxus</i>	L(1, 2)-I	225, 45, 462
	L(3, 4, 5)-II	226, 12, 225, 45, 5, 462
<i>Oeneis chryxus ivallda</i>	L(h?)	5
<i>Neominois ridingsii</i>	L(2[r], 3-4)	5, 545, 462
<i>Lethe portlandia</i>	L	540
<i>Lethe anthedon</i>	L(3-4)	13, 299, 20, 540, 21, 123, 5, 8, 15, 23, 423, 433, 19, 232
<i>Lethe eurydice</i>	L(3-4, 5[r])	12, 13, 299, 474, 20, 22, 433, 21, 15, 5, 8, 19, 130, 144
<i>Lethe appalachia</i>	L(3-4)	481, 474, 295, 9, 540
<i>Lethe creola</i>	L	540
<i>Danaus plexippus</i>	A	19, 20, 21, 519, 8, 299a, 15, 228a, 391, 425, 426, 306a, 306b, 13, 477, 465, 433, 270, 157, 169, 36, 232, 83, 84, 116, 152
<i>Anaea andria</i>	A	13, 299, 402, 540, 277, 23, 276, 5, 273, 339, 340, 151, 332a, 260, 345, 19, 20, 232, 338b, 348d
<i>Nymphalis antiopa</i>	A	12, 427, 13, 284, 540, 236, 243, 529, 341, 342, 344, 345, 369, 276, 394, 235, 8, 215, 207, 21, 23, 177, 94, 87, 78, 303a, 448, 464, 5, 430, 58, 260, 519, 202, 190, 174, 169, 128, 157, 335, 238, 283, 287, 433, 467, 236, 19, 523, 332a, 304, 37, 71, 338b, 349a, 212, 393a
	P(e)	13, 19
<i>Nymphalis milberti</i>	A	427, 13, 337, 394, 369, 529, 433, 15, 283, 23, 62, 87, 78, 335, 332, 19, 177, 129, 169, 348b, 379a, 40b, 413a
	P(e?)	13, 15, 23, 19
<i>Nymphalis californica</i>	A	394, 335, 427, 12, 17, 529, 331b, 422d
	P(e)	17
<i>Nymphalis vau-album</i>	A	19, 21, 343, 23, 327, 15, 524, 433, 13, 299, 12, 156, 78, 87, 349a, 169, 8, 455, 430, 17, 58, 523, 519, 389a, 393a
	P(e)	17
<i>Polygonia interrogationis</i>	A	58, 23, 15, 540, 433, 340, 5, 19, 13, 20, 21, 9, 62, 78, 92, 232, 260, 186, 184, 183, 169, 161, 126, 144, 408a
	P?(e)	19

<i>Polygonia comma</i>	A	58, 519, 13, 299, 482, 340, 276, 346, 433, 9, 125, 78, 92, 5, 540, 23, 15, 260, 169, 262, 19, 20, 21, 37, 232
	P(e)	482
<i>Polygonia progne</i>	A	13, 20, 21, 340, 345, 8, 23, 430, 15, 433, 201, 105, 78, 169, 19, 37
	P(e)	19
<i>Polygonia gracilis</i>	A	19, 52, 212
<i>Polygonia satyrus</i>	A	225, 427, 394, 540, 342, 51, 215
<i>Polygonia zephyrus</i>	A	427, 51, 529, 297a, 331b
<i>Polygonia oreas</i>	A	529
<i>Polygonia faunus</i> (+ hylas)	A	349a, 13, 9, 396, 342, 433, 23, 58, 529, 19, 20, 21, 215, 51, 52
<i>Vanessa atalanta</i>	A	19, 396, 127, 304, 341, 169, 200, 520, 256, 284, 482, 433, 9, 23, 5, 15, 21, 540, 427, 12, 13, 327, 519, 536, 232
	P	19, 482, 9, 23, 15, 21, 540, 12, 13, 519, 348a
<i>Vanessa carye</i>	A	546, 536, 529, 393, 331b
	P	546
	L	546, 331, 520
<i>Vanessa virginiensis</i>	A	331b, 519, 12, 540, 169, 418, 433, 13, 299, 23, 21, 232, 482, 19, 536
	P	519, 12, 433, 13, 299, 23, 481, 15, 21, 482, 19
<i>Vanessa cardui</i>	A	232, 304, 299, 63, 78, 13, 274, 9, 20, 519, 433, 8, 23, 19, 21, 420, 344
	P	13, 423, 20, 519, 23, 19, 21
	L(e?)	331
<i>Limenitis wiedemeyerii</i>	L(3-4)	493, 5
<i>Limenitis lorquini</i>	L(2?, h)	427, 520, 338, 164
<i>Limenitis archippus</i>	L(1[r], 2[r], 3)	23, 13, 299, 21, 74, 103, 131, 455, 5, 8, 460, 459, 160, 232, 393b, 306c, 19, 146, 490, 482, 433, 20, 540, 472, 15, 306, 63, 144, 380a
	P(e)	144
<i>Limenitis arthemis</i>	L(2[r], 3, 4[r])	58, 15, 13, 433, 19, 472, 423, 20, 21, 23, 146, 8, 455, 5, 88, 103, 118, 380a, 393b
<i>L. arthemis astyanax</i>	L(2[r], 3)	13, 299, 540, 472, 23, 19, 20, 21, 121, 132, 430, 433, 393b
<i>Limenitis bredowii</i>	L	520
<i>Asterocampa celtis</i>	L(3-4)	8, 13, 12, 225, 299, 15, 69, 80, 131, 433, 23, 20, 540, 5, 19, 232
	E(e)	12
	A(e?)	225, 433, 20, 5

<i>Asterocampa clyton</i>	L(3-4)	13, 15, 20, 540, 433, 23, 112, 120, 131, 5, 19, 232
<i>A. clyton subpallida</i>	L(3)	516
<i>Boloria titania</i>	L(1, 4)(I&II?)	15, 529, 19, 160, 217, 12
<i>Boloria epithore</i>	L(4)	12
<i>Boloria kriemhild</i>	L(4)	12
<i>Boloria selene</i>	L(2-4)	544, 23, 15, 423, 20, 21, 13, 444, 433, 19, 80, 86, 132, 286a
	L(1)(e)	23, 291, 20, 21, 13, 444, 19, 80
<i>Boloria alberta</i>	L(1)-I	5
	L(h or m) or P-II	5
<i>Boloria eunomia</i>	L(3)	12, 160, 529
<i>Boloria freija</i>	L(4)	529, 160
	L(1)(e)	19
<i>Boloria bellona</i>	L(3-4)	225, 20, 21, 23, 433, 19, 80, 86, 132
	L(1)(e)	225, 13, 15, 21, 23, 19, 80
<i>Speyeria nokomis</i>	L(1, 3[r])	542, 534, 378
<i>Speyeria idalia</i>	L(1)	299, 21, 20, 540, 23, 86, 102, 433, 378
<i>Speyeria cybele</i> (+ <i>leto</i>)	L(1)	299, 19, 20, 21, 540, 109, 75, 86, 5, 120, 132, 15, 180, 524, 23, 72, 12, 181, 397, 433, 37, 378
<i>Speyeria diana</i>	L(1)	9, 299, 540, 5, 23, 19, 72, 75, 86, 109, 120, 132, 378
<i>Speyeria aphrodite</i>	L(1)	19, 20, 21, 433, 109, 107, 23, 5, 423, 15, 413, 72, 75, 86, 378
<i>Speyeria edwardsii</i>	L(1)	12, 150, 378
<i>Speyeria callippe</i>	L(1)	12, 338, 5, 378, 338a
<i>Speyeria atlantis</i>	L(1)	12, 19, 20, 21, 15, 524, 5, 86, 149, 153, 378
<i>Speyeria egleis</i>	L(1)	5, 3, 100, 171, 378
<i>Speyeria zerene</i>	L(1)	44, 542, 378
<i>Speyeria mormonia</i>	L(1)	542, 378
<i>Speyeria coronis</i>	L(1)	137, 462, 542, 378, 338a
<i>Speyeria adiaste</i>	L(1)	542, 378
<i>Speyeria hydaspe</i>	L(1)	542, 378
<i>Dymasia dymas</i>	P(e?)	484
<i>Phyciodes campestris</i>	L(h)	546, 528, 393c
<i>Phyciodes tharos</i>	L(3)	13, 299, 427, 85, 90, 433, 225, 23, 423, 61, 15, 481, 296c, 5, 540, 8, 19, 20, 21, 452, 232, 393c
<i>Phyciodes batesii</i>	L(3)	13, 61, 209, 393c
<i>Phyciodes mylitta</i>	L(h-m)	546, 387
<i>Phyciodes picta</i>	L(2-4)	462, 98, 452, 137, 133
<i>Phyciodes texana</i>	L?	366
<i>Chlosyne theona</i>	L(4)	491, 411
<i>Chlosyne chinatiensis</i>	L(3)	539

<i>Chlosyne leanira</i>	L(3)	529, 427, 338a
<i>Chlosyne leanira alma</i>	L(3)	529
<i>Chlosyne leanira fulvia</i>	L(3)	529, 411
<i>Chlosyne harrisii</i>	L(3-4)	13, 223, 423, 20, 131, 191, 23, 137, 19, 523, 433, 15, 456, 61, 68, 90
<i>Chlosyne nycteis</i>	L(2[r], 3-4)	23, 529, 173, 433, 299, 13, 423, 20, 21, 349, 232, 49, 72, 79, 80, 90, 97, 15, 19
<i>Chlosyne gorgone</i>	L(2, 3, 4)	529, 330, 366, 213, 540, 213, 536, 173, 208
<i>Chlosyne hoffmanni</i>	L(h)	427, 358, 374
<i>Chlosyne palla</i>	L(h)	529, 323
<i>Chlosyne damoetas</i>	L(y)-I?	529
	L(3-4, 5?)-II	529
<i>Chlosyne damoetas malcolmi</i>	L (nearly m)	543a
<i>Chlosyne neuwoegeni</i>	L(h)	427
<i>Chlosyne gabbii</i>	L(3)	50, 338a
<i>Chlosyne californica</i>	L(3)	550, 360
<i>Chlosyne lacinia</i>	L(3)	427, 172, 208, 377d
<i>Poladryas minuta</i>	L(3)	529, 541, 442
<i>P. minuta arachne</i>	L(3)	529
<i>Euphydryas gillettii</i>	L(4-5)	508
<i>Euphydryas chalcedona</i>	L(3[r]-4)	427, 529, 12, 450, 509, 251, 446, 24, 54, 134, 350b
<i>E. chalcedona colon</i>	L(h)	12
<i>E. chalcedona anicia</i>	L(h, 4, 5[r])	529, 167
<i>Euphydryas editha</i>	L(3[r]-4)	138, 154, 137, 427, 98, 439, 480, 462, 296, 98, 479, 5, 350a, 350b
	P(e)	24
<i>Euphydryas phaeton</i>	L(3-4)	13, 299, 327, 76, 79, 540, 90, 5, 134, 137, 15, 165, 462, 37, 393a, 463, 190, 191, 25, 433, 23, 282, 19, 20, 21, 417
Lycaenidae		
<i>Apodemia mormo</i>	L(m)	30, 427, 12, 324, 550
	E(e)	240
<i>Apodemia nais</i>	L(h)	529, 5
<i>A. nais chisosensis</i>	L(almost m)	28
<i>Apodemia palmerii</i>	L(3)	488, 427
<i>Calephelis borealis</i>	L(5-8 of 8-9)	13, 347, 12, 225, 433, 216
<i>Calephelis muticum</i>	L(4-5 of 8-9)	53, 12, 423, 15, 13
<i>Caria ino</i>	L(m)	28
<i>Feniseca tarquinius</i>	P?	19, 20, 21, 13, 142, 163, 433, 232, 481a
	A?	19, 20, 21, 13, 299
	L?	187, 163
<i>Habrodais grunus</i>	E	427, 497

<i>Hypaurotis crysalus</i>	E	353, 529, 34
<i>Phaeostrymon alcestis</i>	E	366, 12
<i>Euristrymon favonius</i>	E	12
<i>E. favonius ontario</i>	E	366, 422
<i>Euristrymon polingi</i>	E	361
<i>Harkenclenus titus</i>	E	13, 299, 12, 19, 20, 21, 15, 433, 466
<i>Satyrium behrii</i>	E	529, 427, 483
<i>Satyrium saepium</i>	E	529, 12, 492
<i>Satyrium californica</i>	E	12, 529
<i>Satyrium tetra</i>	E	427, 12, 500
<i>Satyrium acadica</i>	E	15, 23, 20, 433, 19
<i>Satyrium auretteorum</i>	E	427, 12, 495
<i>Satyrium sylvinus</i>	E	12, 427, 319, 496
<i>Satyrium calanus</i>	E	13, 19, 21, 433, 23, 20, 225, 519
	L(1)(e)	15, 21, 23, 299, 519
	P(e)	122, 19
<i>Satyrium edwardsii</i>	E	20, 476, 13, 433, 15, 19
	L(1)(e)	476, 13, 15, 19
<i>Satyrium liparops</i>	E	12, 13, 19, 20, 423, 433, 15, 466, 23, 58
	L(y)(e)	23
<i>Satyrium caryaevorus</i>	E?	12
<i>Satyrium fuliginosum</i>	E	386, 529, 535
	L?	386
<i>Atlides halesus</i>	P	529, 400, 403, 232
<i>Calycopis cecrops</i>	P?	56
<i>Strymon melinus</i>	P	13, 12, 20, 540, 433, 466, 19, 21, 15, 518, 232
<i>Erora laeta</i>	P?	13, 19, 211
<i>Panthiades m-album</i>	P(e?)	13, 225, 357, 15, 19, 232
<i>Callophrys eryphon</i>	P	427, 352
<i>Callophrys niphon</i>	P	13, 19, 20, 15, 66, 197, 433
<i>Callophrys lanoraieensis</i>	P	523
<i>Callophrys polios</i>	P	13, 198, 433
<i>Callophrys augustinus</i> (* iroides)	P	13, 423, 9, 20, 433, 19, 189, 194, 195
<i>Callophrys mossii</i>	P	552, 530, 529
<i>Callophrys fotis</i>	P	535
<i>Callophrys irus</i>	P	13, 19, 20, 433, 192, 193, 413, 33
<i>Callophrys henrici</i>	P	13, 225, 366, 540, 443, 8, 445, 196
<i>Callophrys spinetorum</i>	P	504
<i>Callophrys gryneus</i>	P	13, 423, 19, 20, 540, 23, 232, 433, 466
<i>Callophrys gryneus nelsoni</i>	P	427, 489
<i>C. gryneus loki</i>	P	427, 550
<i>C. gryneus siva</i>	P	268
<i>Callophrys hesseli</i>	P	57, 413b
<i>Callophrys dumetorum</i>	P	30, 379
	L(e)	471

<i>C. dumetorum viridis</i>	P	253, 379
<i>Callophrys sheridanii</i>	P	382
<i>Callophrys macfarlandi</i>	P	359, 280
<i>Lycaena arota</i>	E	529, 331, 3
<i>Lycaena gorgon</i>	E	427, 494, 470
	L(e)	494, 239
<i>Lycaena helloides</i>	E	529, 363, 546, 368
	L(e)	387
	P(e)	387, 267
<i>Lycaena hyllus</i>	E	13, 20, 437, 15, 19, 433, 529, 203, 60
<i>Lycaena xanthoides</i>	E	427, 500
<i>L. xanthoides editha</i>	E	529
<i>L. xanthoides dione</i>	E	540, 529
<i>Lycaena hermes</i>	E	427, 498, 336, 308
<i>Lycaena epixanthe</i>	E	13, 12, 225, 20, 19, 433, 413, 199, 60, 199
<i>Lycaena dorcas</i>	E	206, 203, 29, 221
<i>Lycaena nivalis</i>	E	544, 310
<i>Lycaena mariposa</i>	E	368
<i>Lycaena heteronea</i>	E	427, 551, 529, 46
<i>Lycaena rubidus</i>	E	551, 529
<i>Lycaena cupreus</i>	L(h)	203, 535
<i>Lycaena phlaeas</i>	L(y, 2-3; m?)	348, 7, 26, 169, 19, 20, 21, 437, 334, 59
	P(e?)	433, 19, 20, 21, 437, 15, 23, 225, 13
<i>Plebejus argyrognomon</i>	E	348, 188, 400
<i>Plebejus melissa</i>	L(h; m?)	43, 19
	P?	19
<i>Plebejus icarioides</i>	L(2)	12, 427, 331, 536, 497, 530, 281, 355, 205
	P(r)	205
<i>Plebejus saepiolus</i>	L(1, h)	12, 13, 427, 513, 214
	P(e?)	438
<i>Plebejus acmon</i>	L(2-3)	333, 382, 383
<i>Plebejus lupini</i>	L(2)	333
<i>Plebejus neurona</i>	L(2)	333
<i>Plebejus emigdionis</i>	L(late larva)	427, 489
<i>Plebejus optilete</i>	L	12
<i>Everes amyntula</i>	L(m)	427, 500, 258, 19, 451
<i>Everes comyntas</i>	L(m)	12, 13, 423, 257, 20, 15, 433, 21, 19
	P(e)	232
<i>Agriades glandon</i>	L or P	529
<i>Brephidium exilis</i>	P	266, 381
<i>Glaucopsyche lygdamus xerces</i>	P	250, 261, 514, 375
<i>Glaucopsyche lygdamus</i>	P	427, 13, 423, 255, 250, 375
	L(m)(e)	19

<i>Glaucoopsyche piasus</i>	P	427, 265
<i>Celastrina argiolus</i>	P	13, 529, 433, 22, 5, 447, 19, 20, 21, 77, 82, 91, 137, 144
	L(e)	433, 144
<i>Celastrina nigra</i> (= <i>ebenina</i>)	P	392
<i>Philotiella speciosa</i>	P	427, 489
<i>Euphilotes rita</i>	P	529
<i>Euphilotes enoptes</i>	P	226, 441
<i>Euphilotes battoides</i>	P	494, 415, 427
Hesperiidae		
<i>Zestusa dorus</i>	P	435
<i>Autochton cellus</i>	P	13
<i>Epargyreus clarus</i>	P	13, 299, 370, 423, 9, 519, 8, 259, 433, 15, 23, 529, 19, 20, 21, 232, 66, 124, 182, 251
<i>Pyrgus communis</i>	L(m)	512, 19, 433, 13, 423
	P(e?)	19, 299, 20, 13, 423
<i>Staphylus hayhurstii</i>	L(3)	309, 322
<i>Pholisora catullus</i>	L(m)	13, 299, 370, 423, 433, 23, 519, 15, 19, 20, 139
	P(r)	9
<i>Erynnis lucilius</i>	L(m)	299, 19, 20, 433
<i>Erynnis persius</i>	L(m)	19, 20, 21, 23, 419, 433
<i>Erynnis baptisiae</i>	L(m)	433
<i>Erynnis brizo</i>	L(m)	9, 19, 20, 21, 66, 433
<i>Erynnis icelus</i>	L(m)	299, 19, 438(h), 18, 136, 433, 20, 21
	P(e)	136
<i>Erynnis juvenalis</i>	L(m)	299, 19, 20, 21, 438h, 372, 9, 23, 66, 522, 519
	P(e)	232
<i>Erynnis zarucco</i>	L	13, 362
<i>Erynnis horatius</i>	L	370
<i>Erynnis tristis</i>	L	488
<i>Erynnis martialis</i>	L(m)	433, 19
<i>Gesta gesta</i>	L(m)	370, 478
<i>Celotes nessus</i>	L	370
<i>Celotes limpia</i>	L	541, 475
<i>Systasea pulverulenta</i>	L(m)	367, 370
<i>Cogia outis</i>	L(m)	370
<i>Cogia caicus</i>	L(m)	27
<i>Thorybes valeriana</i> (= <i>mysie</i>)	L	545
<i>Thorybes pylades</i>	L(m)	492, 499, 427, 433, 178, 370
	P(e)	19, 20, 21, 13
<i>Thorybes bathyllus</i>	L(m)	370, 19, 433
	P(e)	19, 13, 423
<i>Achalarus lyciades</i>	L	370
	P(e)	19, 299, 13

<i>Thymelicus lineola</i>	E	225, 292, 223b, 436, 376a
<i>Copaodes aurantiaca</i>	P?	370
<i>Carterocephalus palaemon</i>	L(m?)	19, 348
<i>Oarisma garita</i>	L(4)	204
<i>Oarisma powesheik</i>	L(5 or 7)	325
<i>Ancyloxypha numitor</i>	L(3-4)	505, 433, 437, 19, 219, 20, 21, 373, 232
	P(e?)	505, 299, 437, 19, 20, 21
<i>Hesperia leonardus</i>	L(1-2)	19, 219, 249, 20, 13, 423, 433
<i>Hesperia leonardus pawnee</i>	L(1)	529, 533a
<i>Hesperia leonardus montana</i>	L(1)	529
<i>Hesperia dacotae</i>	L(5)	533a
<i>Hesperia pahaska</i>	L	521, 427
<i>Hesperia ottoe</i>	L(5)	351, 401
<i>Hesperia metea</i>	L(m)	321, 9, 540, 327
	P(e?)	13, 249, 19
<i>Hesperia columbica</i>	L?	521
<i>Hesperia comma</i>	E	521, 388, 405, 406, 407, 408, 409, 179
<i>Hesperia lindseyi</i>	E	521, 427, 112
<i>Hesperia juba</i>	E or L(1-3)	521
<i>Hesperia nevada</i>	L or P	521
<i>Hesperia miriamae</i>	E?	521
<i>Hesperia sassacus</i>	P?	20, 21, 13, 433, 19
<i>Hesperia viridis</i>	L?	370
<i>Euphyes bimacula</i>	L(h)	529
	P?	19
<i>Euphyes dion</i>	L(3)	540
<i>Euphyes vestris</i>	L(3)	314, 540
<i>Pompeius verna</i>	L?	13, 219
	P?	219
<i>Poanes zabulon</i>	L(m)?	19, 20, 21
	P?	19, 20, 21
<i>Poanes viator</i>	L or P	373
<i>Poanes hobomok</i>	E, L, P?	437, 225, 15, 249, 218, 242, 433
<i>Polites sabuleti</i>	P	317
<i>Polites mardon</i>	P	317
<i>Polites themistocles</i>	P	225, 13, 511, 23, 19, 20, 21, 433, 218
	L?	218
<i>Polites coras</i>	L(1-5)	433, 506, 13, 299, 20, 21, 19
	P	506, 433, 13, 299, 20, 21, 19
<i>Polites origines</i>	L(3-4)	13, 510, 19, 218
	P?	19
<i>Polites mystic</i>	L(4)	226, 423, 507, 19, 21, 20, 13, 433
<i>Ochlodes sylvanoides</i>	L(1)	331, 427
	E?	427
<i>Paratrytone melane</i>	L?	486

<i>Problema byssus</i>	L(4)	371, 12, 540
<i>Atrytone logan</i>	L?	13, 423, 19
	P?	13, 423
<i>Atrytone arogos</i>	L(4 of 6)	327, 540, 481
<i>Panoquina panoquinoides errans</i>	L	485
<i>Wallengrenia egeremet</i>	L(h)	13, 423, 433, 19, 218, 249
<i>Amblyscirtes hegon</i>	L(1)	399
	P?	19
<i>Amblyscirtes celia belli</i>	L(4)	316
<i>Amblyscirtes nysa</i>	L(h-m)	315, 540
<i>Amblyscirtes aenus</i>	L(h)	529
<i>A. aenus linda</i>	L	540
<i>Amblyscirtes vialis</i>	L	540
	P(e)	19, 13, 423, 23
<i>Atrytonopsis hianna</i>	L(m)	329, 540
<i>Megathymus yuccae</i>	L(h-m)	12, 495, 39, 25, 326, 517, 25b, 25c
<i>Megathymus ursus</i>	L(3, 4, 5)	12, 25b
<i>M. ursus violae</i>	L(h)	529
<i>Megathymus cofaqui</i>	L	263
<i>M. cofaqui harrisi</i>	L	9
<i>Megathymus streckeri texanus</i>	L	271
<i>Megathymus streckeri streckeri</i>	L	26a
<i>Stallingsia smithi maculosus</i>	L	350
<i>Agathymus neumogeni</i>	L(1-3)	311, 38
<i>Agathymus polingi</i>	L(1-3)	311
<i>Agathymus evansi</i>	L(1-3)	311
<i>Agathymus aryxna</i>	L(1-3)	311, 12
<i>A. aryxna baueri</i>	L(1-3)	311
<i>A. aryxna freemani</i>	L(1-3)	311
<i>Agathymus alliae</i>	L(1-3)	311, 12
<i>Agathymus mariae</i>	L	313, 224

Table 2

Species which probably have no diapause stage. These are mainly subtropical species or migrants to temperate areas. For explanation of symbols see Table 1. n-source states that no diapause occurs.

SPECIES	WINTER STAGE	SOURCE
<i>Papilio palamedes</i>	P	13, 23, 366(n), 19, 114
	L	13, 23
<i>Nathalis iole</i>	A	540, 296b(n)
	P	233
<i>Phoebis sennae</i>	A	15, 23, 21, 20, 19
	P?	232
	L?	21, 20, 19
<i>Phoebis agarithe</i>	L(y)	297
<i>Eurema lisa</i>	A	15(n), 23, 12, 13, 482(n), 20, 19(n)
	P	12, 13, 232
<i>Colias cesonia</i>	A	327(n), 273, 404, 277, 540
	P	13, 427, 232, 363a, 370a(n)
<i>Ascia monuste</i>	L, P, A	302, 47(n)
<i>Euptychia hermes</i>	L	320, 366(n), 13, 232
<i>Agraulis vanillae</i>	L, P, A	332a(n), 395(n), 288(n), 232, 364a, 260a(n), 370a(n)
<i>Euptoieta claudia</i>	A	23, 423, 540, 21, 13, 225, 12, 20, 19, 232, 277
	P	23, 540, 21, 20, 19
	L	23, 8, 21, 20, 19, 16, 277
<i>Precis coenia</i>	A	232, 482, 23, 13, 12, 540, 433, 327(n), 520(n), 19, 20, 21
	P?	546, 387
	L	546, 529, 387
<i>Calephelis nemesis</i>	L or P	487
<i>Ministrymon leda</i>	P?	501
<i>Strymon columella</i>	L?	498
<i>Panoquina ocola</i>	L or P	364
<i>Nyctelius nyctelius</i>	L	301
<i>Lerodea eufala</i>	L, P?	264
<i>Polites vibex</i>	L or P	370
<i>Atalopedes campestris</i>	A?	13
<i>Lerema accius</i>	P?	13, 19
<i>Perichares philetas</i>	L	301
<i>Calpodas ethlius</i>	L or P	370, 482(n), 260a(n)
<i>Urbanus proteus</i>	A	225, 19
<i>Helioptetes macaira</i>	?	370(n)
<i>Helioptetes laviana</i>	L	370
<i>Phocides polybius</i>	L	356
<i>Cogia calchas</i>	L	372

<i>Timochares ruptifasciatus</i>	L	389
<i>Cabares potrillo</i>	L	389
<i>Xenophanes trixus</i>	L	389
<i>Chioides catillus</i>	L	372, 370
<i>Staphylus mazans</i>	L or P	370
<i>Chiomara asychis</i>	?	389(n)

Table 3

Hibernal diapause stage of Nearctic, Japanese, and British butterflies (E-egg, L-larva, P-pupa, A-adult).

	Nearctic				Japan				Britain			
	E	L	P	A	E	L	P	A	E	L	P	A
1. Papilionidae												
<i>Parnassius</i>	3	2	1		3		1					
other Papilionidae			17				13				1	
2. Pieridae												
Coliadinae		10		1		2	1	4		2	1	1
Pierinae	1		13			3	5				3	
3. Lycaenidae												
Theclini	17		21		25	1	2	3	4			1
Miletini						1						
Curetini								1				
Lycaenini	12	2				1					2	
Polyommataini	1	10	10		4	8	4		2	9	1	
Riodininae		6										1
4. Nymphalidae												
<i>Polygonia, Nymphalis,</i>												
<i>Aglais, Vanessa</i>				12				10				6
<i>Anaea</i>				1								
Danainae				1								
Satyrinae		27				18	1	1		10		
other Nymphalidae		52	1		2	23	3	1	1	8		
5. HesperIIDae	3	59	7		2	13	2		2	5	1	
Total	37	168	70	15	36	70	32	20	9	36	9	7

Table 4

Biennial species: hibernal diapause stage, cohort of adult flight (even or odd years), and documentation of biennialism. Diapause stages from Table 1.

SPECIES	HIBERNAL DIAPAUSE STAGE	ADULT FLIGHT TIME & LOCATION	DOCUMENTATION FOR BIENNIALISM
<i>Parnassius eversmanni</i>	E-I P-II	Japan	348
<i>Parnassius phoebus, clodius</i>	E-I, L-II	----	----
<i>Colias meadii</i>	L(1)-I L(3-4, 5 [r])-II	Alberta	162
<i>Colias nastes</i>	L(y,3,m)- (I+II?) P?	----	----
<i>Colias hecla</i>	L(I+II), P?	----	12, 13
<i>Chlosyne damoetas</i>	L(y)-I? L(3-4, 5?)- II	every yr. Colo.	529
<i>Boloria alberta</i>	L(1)-I L(h or m) or P-II	mostly even yrs. Alberta, mostly odd yrs. Plateau Mtn. Alberta	5, 200, 247, 529, 398a 414b, 12
<i>Boloria astarte</i>	----	even yrs. Washington, even and odd yrs. Alberta	414b, 12, 17, 529, 398a
<i>B. astarte distincta</i>	----	mostly even yrs. in north, mostly odd yrs. in south	*, 370b
<i>Boloria polaris</i>	----	odd yrs. central Alaska, Manitoba, mostly even yrs. rest of range	*, 385, 329a, 393d, 422c, 414c, 331a
<i>B. (titania?) chariclea</i>	L(1,4) (titania)	odd yrs. Alaska, Yukon	*, 370b
<i>Oeneis jutta</i>	L(1, 2, 3)-I L(4, 5, 6)-II	odd yrs. Alaska, Michigan, Wisconsin, Minnesota, Sask- atchewan; even yrs. Colo., Manitoba, Newfoundland, Ontario, New Brunswick, Maine, Labrador, Quebec	385, 42, 168, 25a, 419a, 417a, 398b, 414c, 529, 10, *, 377c, 422c, 422a, 414a, 329a, 331a
<i>Oeneis polixenes</i>	L(1, per- haps 2- 3)-I L(4, 5)-II	odd yrs. Alaska; even yrs. Maine	*, 351a, 422c

<i>Oeneis melissa</i>	L(y, 2, 3)-I L(5)-II	every yr. Colo.	167, 5, 19
<i>Oeneis uhleri</i>	L(2, 3, 4, 5)(I+II?)	biennial Alberta; may be more common odd yrs. Colo.	5, 529
<i>Oeneis nevadensis</i>	L(2,3)-I L(5)-II	even yrs. Wash., Vancouver Island; mostly even yrs. Oregon & Sonoma Co. Calif.	395a, 5, 369, 17, 385, 398, 332, 422e, 12, 344a, 300
<i>Oeneis (nevadensis?) macounii</i>	L(1, 2)-I L(m)-II	odd yrs. Alberta, Saskatchewan, Riding Mts. Manitoba; even yrs. eastern Manitoba, Ontario, Minnesota	167, 385, 10, 374a, 422a, 398c, 168, 175, 375a
<i>Oeneis chryxus</i>	L(1, 2)-I L(3,4,5)-II	even yrs. Michigan & so. Colo., odd yrs. Sonora Pass Calif.; mostly odd rest of Calif. (<i>ivallda</i>)	385, 529, 307, 416, 422e
<i>Oeneis taygete</i>	-----	odd- & even-yr. cohorts differ in phenotype in N. Wyoming; even yrs. Labrador	385, 537, 414c, 543, 414
<i>Oeneis alberta</i>	-----	biennial Colorado?	553
<i>Neominois ridingsii</i>	L(2[r], 3-4)+?	even yrs. Sonora Pass Calif.	529
<i>Erebia disa</i>	-----	odd yrs. Alaska, Saskatchewan; even yrs. part of Saskatchewan	*, 385, 10, 318a, 422c
<i>Erebia theano</i>	L(y)+?-I&II	even yrs. throughout Colo.	385, 529
<i>Hesperia comma</i>	E + ?	odd yrs. Alaska; even yrs. Churchill Manitoba	*, 329a
<i>Pyrgus centaureae</i>	-----	mostly odd yrs. Colo.	529

*references 405, 406, 407, 408, 409

